


Any Disorder

Prevalence


- **12-month Prevalence:** 26.2% of U.S. adult population¹
- **Severe:** 22.3% of these cases (e.g., 5.8% U.S. adult population) are classified as “severe”²



Average Age-of-Onset: 14 years old⁴


Demographics (for lifetime prevalence)⁵

- **Sex:** Women are no more or less likely than men to experience any disorder over their lifetime
- **Race:** Non-Hispanic blacks are 30% less likely than non-Hispanic whites to experience any disorder during their lifetime
- **Age:**




Treatment/Services Use⁶

- **12-month Healthcare Use:** 36.0% of those with disorder are receiving treatment
- **Percent Received Minimally Adequate Treatment:** 33.4% of those receiving treatment are receiving minimally adequate treatment (12.0% of those with disorder)



- **12-month Any Service Use (including Healthcare):** 41.1% of those with disorder are receiving treatment
- **Percent Received Minimally Adequate Treatment:** 32.7% of those receiving treatment are receiving minimally adequate treatment (13.4% of those with disorder)



¹Kessler RC, Chiu WT, Demler O, Walters EE. Prevalence, severity, and comorbidity of twelve-month DSM-IV disorders in the National Comorbidity Survey Replication (NCS-R). *Archives of General Psychiatry*. 2005 Jun;62(6):617-27.

²Ibid

³Kessler RC, Berglund PA, Demler O, Jin R, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication (NCS-R). *Archives of General Psychiatry*. 2005 Jun;62(6):593-602.

⁴Ibid

⁵Ibid

⁶Wang PS, Lane M, Olfson M, Pincus HA, Wells KB, Kessler RC. Twelve month use of mental health services in the United States. *Archives of General Psychiatry*. 2005 Jun;62(6):629-640.